

**Comparison of antidepressant and side effects in mice after
intranasal administration of (*R,S*)-ketamine, (*R*)-ketamine,
and (*S*)-ketamine**

(RS-ケタミン、R-ケタミンおよびS-ケタミンの鼻腔内投与後の抗うつ作用
および副作用の比較)

千葉大学大学院医学薬学府

先端医学薬学専攻

(主任:橋本謙二教授)

常 立甲

TABLE OF CONTENTS

Abstract	3
Introduction	4
Materials and methods	7
Results	12
Discussion	19
Acknowledgments	22
References	23

ABSTRACT

The *N*-methyl-D-aspartate receptor (NMDAR) antagonist (*R,S*)-ketamine produces rapid and sustained antidepressant effects in treatment-resistant patients with depression although intranasal use of (*R,S*)-ketamine in ketamine abusers is popular. In March 5, 2019, nasal spray of (*S*)-ketamine for treatment-resistant depression was approved as a new antidepressant by the US Food Drug Administration. Clinical study of (*R*)-ketamine is underway. In a chronic social defeat stress (CSDS) model, we compared the antidepressant effects of (*R,S*)-ketamine, (*R*)-ketamine, and (*S*)-ketamine after a single intranasal administration. Furthermore, we also compared the side effects (i.e., locomotion, prepulse inhibition (PPI), abuse liability) of these three compounds in mice. The order of potency of antidepressant effects after a single intranasal administration was (*R*)-ketamine > (*R,S*)-ketamine > (*S*)-ketamine. In contrast, the order of locomotor activity and prepulse inhibition (PPI) deficits after a single intranasal administration was (*S*)-ketamine > (*R,S*)-ketamine > (*R*)-ketamine. In the conditioned place preference (CPP) test, both (*S*)-ketamine and (*R,S*)-ketamine increased CPP scores in mice after repeated intranasal administration, in a dose dependent manner. In contrast, (*R*)-ketamine did not increase CPP scores in mice. These findings suggest that intranasal administration of (*R*)-ketamine would be a safer antidepressant than (*R,S*)-ketamine and (*S*)-ketamine.

Keywords: Antidepressant; (*R*)-Ketamine; (*R,S*)-Ketamine; (*S*)-Ketamine; Side effects.

Introduction

In 2010, Berman et al. (2000) reported a first double-blind, placebo-controlled study of the N-methyl-D-aspartate receptor (NMDAR) antagonist (*R,S*)-ketamine, demonstrating that (*R,S*)-ketamine exhibits rapid antidepressant effects in treatment-resistant patients with major depressive disorder (MDD). Subsequently, a number of groups replicated robust antidepressant effects of (*R,S*)-ketamine in treatment-resistant patients with MDD (Murrough et al., 2013; Su et al., 2017; Zarate et al., 2006;). Interestingly, (*R,S*)-ketamine could produce anti-suicidal effects in treatment-resistant patients with MDD (Grunebaum et al., 2018; Larkin et al., 2011; Murrough et al., 2015; Price et al., 2009). Several meta-analyses showed that (*R,S*)-ketamine exhibits rapid antidepressant and anti-suicidal ideation effects in treatment-resistant patients with MDD or bipolar disorder (Kishimoto et al., 2016; Newport et al., 2015; Wilkinson et al., 2018; Xu et al., 2016). Off-label use of (*R,S*)-ketamine (i.e., intravenous and intranasal administration) for antidepressant effects is increasing in the United State of America (USA) although the common adverse effects (e.g., psychotomimetic effects and dissociative effects) of (*R,S*)-ketamine are not resolved (Singh et al., 2017; Wilkinson et al., 2017; Zhu et al., 2017). Thus, although (*R,S*)-ketamine is the most attractive antidepressant in the treatment of severe depression, the precise mechanisms underlying its antidepressant actions remain elusive (Abdallah et al., 2018; Chaki, 2017a; 2017b; Duman, 2018; Gould et al., 2019; Hashimoto, 2016a; 2016b; Krystal et al., 2019; Monteggia and Zarate, 2015; Murrough et al., 2017; Zanos et al., 2018; Zhang and Hashimoto, 2019a).

(*R,S*)-ketamine ($K_i = 0.53 \mu\text{M}$ for NMDAR) is a racemic mixture containing equal parts of (*R*)-ketamine (or arketamine) ($K_i = 1.4 \mu\text{M}$ for NMDAR) and (*S*)-ketamine (or esketamine) ($K_i = 0.30 \mu\text{M}$ for NMDAR) (Ebert et al., 1997). (*R*)-ketamine is reported to show greater potency and longer-lasting antidepressant effects than (*S*)-ketamine in several animal models of depression (Fukumoto et al., 2017; Yang et al., 2015; 2017a; 2017b; 2018a;

Zanos et al., 2016; Zhang et al., 2014). Unlike (*S*)-ketamine, (*R*)-ketamine might not induce psychotomimetic side effects or exhibit abuse potential in rodents (Yang et al., 2015; 2016). In addition, unlike (*R,S*)-ketamine and (*S*)-ketamine, (*R*)-ketamine did not cause the expression of heat shock protein HSP-70 (a marker for neuronal injury) in the rat retrosplenial cortex after a single intraperitoneal (i.p.) administration (Tian et al., 2018). A positron emission tomography (PET) study showed a marked reduction of dopamine D2/3 receptor binding in conscious monkey striatum after a single intravenous (i.v.) infusion of (*S*)-ketamine but not that of (*R*)-ketamine, suggesting that (*S*)-ketamine-induced dopamine release might be associated with acute psychotomimetic and dissociative side effects in humans (Hashimoto et al., 2017). Therefore, it seems that (*R*)-ketamine could be a safer antidepressant in humans than (*R,S*)-ketamine and (*S*)-ketamine (Hashimoto, 2014; 2016a; 2016b; 2016c).

Fukumoto et al. (2017) reported that (*R,S*)-ketamine (10 mg/kg) and (*R*)-ketamine (10 mg/kg), but not (*S*)-ketamine (3 and 10 mg/kg), significantly reversed the depressive-like behavior induced by repeated treatments with corticosterone in rats at 24 hours after a single i.p. administration, indicating that (*S*)-ketamine's antidepressant effects are less potent than (*R,S*)-ketamine and (*R*)-ketamine. On March 5, 2019, the US Food Drug Administration (FDA) approved Janssen Pharmaceutical Inc.'s (*S*)-ketamine nasal spray for treatment-resistant depression (FDA 2019). It is well known that bioavailability (17 – 29%) of intranasal administration of (*R,S*)-ketamine in humans is markedly lower than i.v. (100%) and intramuscular (i.m.) administration (93%) (Li et al., 2016; Pelltoniemi et al., 2016; Zhang and Hashimoto, 2019a), suggesting lower efficacy and higher individual difference of intranasal administration compared to i.v. and i.m. administration. However, there are no reports showing the direct comparison of intranasal administration of (*R,S*)-ketamine and its two enantiomers for antidepressant and side effects in rodents.

The purpose of this study is to compare the antidepressant and side effects of intranasal administration of (*R,S*)-ketamine and its two enantiomers (*R*)-ketamine and (*S*)-ketamine. First, we compared the antidepressant effects of a single intranasal administration of (*R,S*)-ketamine, (*R*)-ketamine and (*S*)-ketamine in susceptible mice after chronic social defeat stress (CSDS). Second, we compared the side effects [i.e., locomotion, prepulse inhibition (PPI), conditioned place preference (CPP)] of intranasal administration of (*R,S*)-ketamine, (*R*)-ketamine and (*S*)-ketamine in mice.

Material and methods

Animals

Male adult C57BL/6 mice (n = 400), aged 8 weeks (body weight 20-25g, Japan SLC, Inc., Hamamatsu, Japan) and male adult CD1 (ICR) mice (n = 40), aged 13-15 weeks (body weight >40g, Japan SLC, Inc., Hamamatsu, Japan) were used. Animals were housed under controlled temperatures and 12-hour light/dark cycles (lights on between 07:00-19:00 h), with ad libitum food (CE-2; CLEA Japan, Inc., Tokyo, Japan) and water. The protocol was approved by the Chiba University Institutional Animal Care and Use Committee (Permission number: 29-420). This study was carried out in strict accordance with the recommendations in the Guide for the Care and Use of Laboratory Animals of the National Institutes of Health, USA. Animals were deeply anaesthetized with isoflurane before being killed by cervical dislocation. All efforts were made to minimize suffering.

Materials

(*R*)-ketamine hydrochloride and (*S*)-ketamine hydrochloride were prepared by recrystallization of (*R,S*)-ketamine (Ketalar®, ketamine hydrochloride, Daiichi Sankyo Pharmaceutical Ltd., Tokyo, Japan) and D-(-)-tartaric acid and L-(+)-tartaric acid, respectively (Zhang et al., 2014). The dose (10, 20 or 40 mg/kg as hydrochloride) of (*R,S*)-ketamine and its enantiomers dissolved in the physiological saline was used as previously reported (Chang et al., 2019; Yang et al., 2015; 2017a; 2017b; 2018a; Zhang et al., 2018). Other reagents were purchased commercially.

Chronic social defeat stress (CSDS) model

The procedure of CSDS was performed as previously reported (Chang et al., 2019; Dong et al., 2017; Golden et al., 2011; Yang et al., 2015; 2017a; 2017b; 2018a; Xiong et al., 2018a; 2018b; Zhang et al., 2018). The C57BL/6 mice were exposed to a different CD1 aggressor mouse for 10 min per day for consecutive 10 days. When the social defeat session ended, the

resident CD1 mouse and the intruder mouse were housed in one half of the cage separated by a perforated Plexiglas divider to allow visual, olfactory, and auditory contact for the remainder of the 24-h period. At 24 h after the last session, all mice were housed individually. On day 11, a social interaction test (SIT) was performed to identify subgroups of mice that were susceptible and unsusceptible to social defeat stress. This was accomplished by placing mice in an interaction test box (42×42 cm) with an empty wire-mesh cage (10×4.5 cm) located at one end. The movement of the mice was tracked for 2.5 min, followed by 2.5 min in the presence of an unfamiliar aggressor confined in the wire-mesh cage. The duration of the subject's presence in the "interaction zone" (defined as the 8-cm-wide area surrounding the wiremesh cage) was recorded by a stopwatch. The interaction ratio was calculated as time spent in an interaction zone with an aggressor/time spent in an interaction zone without an aggressor. An interaction ratio of 1 was set as the cutoff: mice with scores < 1 were defined as "susceptible" to social defeat stress and those with scores \geq 1 were defined as "resilient". Approximately 70-80 % of mice were susceptible after CSDS. Susceptible mice were randomly divided in the subsequent experiments. Control C57BL/6 mice without CSDS were housed in the cage before the behavioral tests.

Treatment and behavioral tests

The CSDS susceptible mice were divided to four groups. Subsequently, saline (0.5 ml/kg), (*R,S*)-ketamine (10 mg/kg), (*R*)-ketamine (10 mg/kg), or (*S*)-ketamine (10 mg/kg) was administered intranasally into CSDS susceptible mice (Figure 1A). Mice were restrained by hand, and saline or ketamine was administered intranasally into awake mice using Eppendorf micropipette (Eppendorf Japan, Tokyo, Japan). Behavioral tests, including locomotion test (LMT), tail suspension test (TST), forced swimming test (FST) and 1% sucrose preference test (SPT), were performed as reported previously (Dong et al., 2017; Yang et al., 2015; 2017a; 2017b; 2018a; Xiong et al., 2018a; 2018b; Zhang et al., 2018). LMT and TST were

performed 2 and 4 hours after a single injection, respectively. FST was performed 1 day after injection. SPT was performed 2, and 7 days after a single injection (Figure 1A).

Locomotion

The locomotor activity was measured by an animal movement analysis system SCANETMV-40 (MELQUEST Co., Ltd., Toyama, Japan). The mice were placed in experimental cages (length× width × height: 560 × 560 × 330 mm). The cumulative locomotor activity counts were recorded for 60 minutes. Cages were cleaned between testing session.

Tail suspension test (TST)

A small piece of adhesive tape placed approximately 2 cm from the tip of the tail for mouse. A single hole was punched in the tape and mice were hung individually, on a hook. The immobility time was recorded for 10 minutes. Mice were considered immobile only when they hung passively and completely motionless.

Forced swimming test (FST)

The FST was conducted using an automated forced-swim apparatus (SCANET MV-40; MELQUEST Co., Ltd., Toyama, Japan). Mice were placed individually in a cylinder (diameter: 23 cm; height: 31 cm) containing 15 cm of water maintained at a temperature of 23°C ± 1°C. The immobility time was calculated using the activity time as (total) – (active) time by the apparatus analysis software. The immobility time of each mouse was recorded for a period of 6 min.

Sucrose preference test (SPT)

Mice were exposed to water and 1% sucrose solution for 48 h, followed by 4 hours of water and food deprivation and a 1- hour exposure to two identical bottles (water and 1% sucrose solution). The bottles containing water and sucrose were weighed before and at the end of this period. The sucrose preference was calculated as a percentage of sucrose solution consumption to the total liquid consumption.

Side effects

Locomotion

After habituation (60 min) in the cage, saline (0.5 ml/kg), (*R,S*)-ketamine (10, 20 or 40 mg/kg), (*R*)-ketamine (10, 20 or 40 mg/kg), or (*S*)-ketamine (10, 20 or 40 mg/kg) was injected intranasally into male C57BL/6 mice. Locomotor activity was measured using an animal movement analysis system (SCANET MV-40, Melquest, Toyama, Japan). The system consisted of a rectangular enclosure (560 x 560 mm). The side walls (height, 60 mm) of the enclosure were equipped with 144 pairs of photosensors located at 6-mm intervals at a height of 30 mm from the bottom edge. An animal was placed in the observation cage 60 minutes from a single dose of saline or compounds. A pair of photosensors was scanned every 0.1 second to detect the animal's movements. The intersection of paired photosensors (10 mm apart) in the enclosure was counted as one unit of locomotor activity. Data collected for 60 minutes after a single injection were used in this study.

Pupulse inhibition (PPI) test

Male C57BL/6 mice were tested for their acoustic startle reactivity (ASR) in a startle chamber (SR-LAB; San Diego Instruments, San Diego, CA, USA) using the standard methods described previously (Matsuura et al., 2015; Yang et al., 2015; Yang et al., 2018b). The test sessions were begun after an initial 10-min acclimation period in the chamber. The mice were subjected to one of six trials: (1) pulse alone, as a 40 ms broadband burst; a pulse (40 ms broadband burst) preceded by 100 ms with a 20 ms prepulse that was (2) 4 dB, (3) 8 dB, (4) 12 dB, or (5) 16 dB over background (65 dB); and (6) background only (no stimulus). The amount of PPI was expressed as the percentage decrease in the amplitude of the startle reactivity caused by presentation of the prepulse (% PPI). Saline (0.5 ml/kg), or (*R,S*)-ketamine (10, 20 or 40 mg/kg) [or (*R*)-ketamine (10, 20 or 40 mg/kg), (*S*)-ketamine (10, 20 or

40 mg/kg)] was administered intranasally 20 min (including the 10-min acclimation period) before the machine records. The PPI test lasted 20 min in total.

Conditioned place preference (CPP) test

The place conditioning paradigm (Brain Science Idea Inc., Osaka, Japan) was used for studying ketamine-induced rewarding effects, as reported previously (Yang et al., 2015; Yang et al., 2018b). Male C57BL/6 mouse was allowed to move freely between transparent and black boxes for a 15 min session once a day, for 3 days (days 1-3) as preconditioning. On day 3, the time spent in each box was measured. There was no significant difference between time spent in the black compartment with a smooth floor and the white compartment with a textured floor, indicating that there was no place preference before conditioning. On days 4, 6, and 8, saline (0.5 ml/kg), or (*R,S*)-ketamine (10, 20 or 40 mg/kg) [or (*R*)-ketamine (10, 20 or 40 mg/kg), (*S*)-ketamine (10, 20 or 40 mg/kg)] was intranasally administered, and then mice were confined to either the transparent or black box for 30 min. On days 5, 7, and 9, mice were given saline and placed in the opposite ketamine-conditioning box for 30 min. On day 10, the post-conditioning test was performed without drug treatment, and the time spent in each box was measured for 15 min. A counterbalanced protocol was used in order to nullify any initial preference by the mouse. The CPP score was designated as the time spent in the drug-conditioning sites, minus the time spent in the saline-conditioning sites.

Statistical analysis

The data show as the mean \pm standard error of the mean (S.E.M.). Analysis was performed using PASW Statistics 20 (formerly SPSS Statistics; SPSS, Tokyo, Japan). The data were analyzed using the one-way analysis of variance (ANOVA), followed by post-hoc Fisher's Least Significant Difference (LSD) test. The PPI data were also analyzed using multivariate analysis of variance, followed by post-hoc Fisher's LSD test. The *P*-values of less than 0.05 were considered statistically significant.

Results

Antidepressant effects of (*R,S*)-ketamine, (*R*)-ketamine and (*S*)-ketamine in CSDS susceptible mice

Locomotion showed no difference ($F_{4,72} = 0.735$, $P = 0.571$) among the five groups (Figure 1B). One-way ANOVA of TST data showed a statistical significance ($F_{4,72} = 15.23$, $P < 0.001$) among the five groups (Figure 1C). Post-hoc tests showed that (*R,S*)-ketamine (10 mg/kg) and (*R*)-ketamine (10 mg/kg) significantly attenuated the increased immobility times of TST in CSDS susceptible mice (Figure 1C). However, (*S*)-ketamine (10 mg/kg) did not attenuate the increased immobility time of TST in CSDS susceptible mice although (*S*)-ketamine slightly decreased the increased immobility time (Figure 1C). One-way ANOVA of FST data showed a statistical significance ($F_{4,72} = 13.77$, $P < 0.001$) among the five groups (Figure 1D). Post-hoc tests showed that three compounds (10 mg/kg) significantly attenuated the increased immobility times of FST in CSDS susceptible mice (Figure 1D). One-way ANOVA of SPT data showed statistical significance (2 days after a single injection: $F_{4,72} = 20.78$, $P < 0.001$) among the five groups (Figure 1E). Post-hoc tests showed that three compounds (10 mg/kg) significantly attenuated the decreased sucrose preference of SPT in CSDS susceptible mice (Figure 1E). One-way ANOVA of SPT data showed statistical significance (7 days after a single injection: $F_{4,72} = 9.311$, $P < 0.001$,) among the five groups (Figure 1F). Post-hoc tests showed that sucrose preference of (*R*)-ketamine-treated group was significantly higher from saline-treated group. However, sucrose preference of (*R,S*)-ketamine-treated group and (*S*)-ketamine -treated group was not different from saline-treated group (Figure 1E and 1F). Collectively, the order of potency of antidepressant effects in a CSDS model was (*R*)-ketamine > (*R,S*)-ketamine > (*S*)-ketamine.




Figure 1. Schedule of a CSDS model, treatment, and behavioral tests

(A): CSDS was performed from day 1 to day 10, and the social interaction test (SIT) was performed on day 11. Saline (0.5 ml/kg), (*R,S*)-ketamine (10 mg/kg), (*R*)-ketamine (10 mg/kg), or (*S*)-ketamine (10 mg/kg) was administered intranasally into the susceptible mice on day 12. LMT and TST were performed 2 and 4 hours after a single injection, respectively. SPT was performed 2, and 7 days after a single injection. (B): LMT. (day 12). (C): TST (day 12). (D): FST (day 13). (E): SPT (day 14). (F): SPT (day 19). The values represent the mean \pm S.E.M. ($n = 15$ or 16). $*P < 0.05$, $**P < 0.01$, $***P < 0.001$ compared with saline-treated susceptible mice. N.S.: not significant. LMT: locomotion test. TST: tail suspension test. FST: forced swimming test. SPT: 1% sucrose preference test. R-KT: (*R*)-ketamine. RS-KT: (*R,S*)-ketamine. S-KT: (*S*)-ketamine.

Effects of (*R,S*)-ketamine, (*R*)-ketamine, and (*S*)-ketamine on locomotion in mice after a single intranasal administration.

Effects of three compounds on locomotion of male mice were examined after a single intranasal administration. One-way ANOVA of the data showed statistical significances ($F_{9,70} = 8.931, P < 0.001$) among the ten groups (Figure 2). Post-hoc tests showed that a single intranasal administration of (*R,S*)-ketamine (20 and 40 mg/kg) or (*S*)-ketamine (10, 20 and 40 mg/kg) significantly increased locomotion compared to saline-treated group (Figure 2). Furthermore, locomotion of (*S*)-ketamine (40 mg/kg) treated mice was significantly higher than that of (*R,S*)-ketamine (40 mg/kg) or (*R*)-ketamine (40 mg/kg) treated mice. In contrast, all doses (10, 20 or 40 mg/kg) of (*R*)-ketamine did not alter locomotion in mice (Figure 2).




Figure 2. Effects of (*R,S*)-ketamine, (*R*)-ketamine, and (*S*)-ketamine on locomotion after a single intranasal administration

Saline (0.5 ml/kg), (*R,S*)-ketamine (10, 20, or 40 mg/kg), (*R*)-ketamine (10, 20, or 40 mg/kg), or (*S*)-ketamine (10, 20, or 40 mg/kg) was administered intranasally into male mice. Locomotor activity was measured 60 min after a single injection of the compounds. The values represent the mean \pm S.E.M. ($n = 8$). * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$ compared with saline-treated mice. & $P < 0.05$ compared to RS-KT (40 mg/kg). &&& $P < 0.001$ compared to R-KT (40 mg/kg). N.S.: not significant. R-KT: (*R*)-ketamine. RS-KT: (*R,S*)-ketamine. S-KT: (*S*)-ketamine.

Effects of (*R,S*)-ketamine, (*R*)-ketamine, and (*S*)-ketamine on PPI in mice after a single intranasal administration

PPI test was performed to examine the effects of three compounds in mice. There were no changes in the acoustic startle response among the four groups for (*R,S*)-ketamine [Wilks lambda = 0.717, $P = 0.589$], (*R*)-ketamine [Wilks lambda = 0.629, $P = 0.226$], and (*S*)-ketamine [Wilks lambda = 0.588, $P = 0.405$] (Figure 3A-3C).

The MANOVA analysis of all PPI data of (*R,S*)-ketamine revealed that there was a significant effect [Wilks lambda = 0.497, $P = 0.042$]. Treatment with (*R,S*)-ketamine (10, 20 or 40 mg/kg) decreased PPI at all dB groups, in a dose dependent manner. Subsequent post-hoc tests indicated significant differences in PPI between the saline group and (*R,S*)-ketamine (40 mg/kg) group at all dB groups (Figure 3A). In contrast, the MANOVA analysis of all PPI data of (*R*)-ketamine revealed that there was not a significant effect [Wilks lambda = 0.625, $P = 0.216$] (Figure 3B). The MANOVA analysis of all PPI data of (*S*)-ketamine revealed that there was a significant effect [Wilks lambda = 0.299, $P < 0.001$]. Treatment with (*S*)-ketamine (10, 20 or 40 mg/kg) decreased PPI at all dB groups, in a dose dependent manner. Subsequent post-hoc tests indicated significant differences in PPI deficits between the saline group and (*S*)-ketamine (20 and 40 mg/kg) group at all dB groups. Furthermore, (*S*)-ketamine (10 mg/kg) significantly decreased PPI at 73 dB (Figure 3C).




Figure 3. Effects of (*R,S*)-ketamine, (*R*)-ketamine, and (*S*)-ketamine on PPI after a single intranasal administration

(A): Saline (0.5 ml/kg), or (*R,S*)-ketamine (10, 20, or 40 mg/kg) was administered intranasally into male mice. (B): Saline (0.5 ml/kg), or (*R*)-ketamine (10, 20, or 40 mg/kg) was administered intranasally into male mice. (C): Saline (0.5 ml/kg), or (*S*)-ketamine (10, 20, or 40 mg/kg) was administered intranasally into male mice. Startle response amplitude and PPI were measured as described in the method section. The values represent the mean \pm S.E.M. (n = 9 or 10). * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$ compared with saline-treated mice. N.S.: not significant. R-KT: (*R*)-ketamine. RS-KT: (*R,S*)-ketamine. S-KT: (*S*)-ketamine.

Effects of (*R,S*)-ketamine, (*R*)-ketamine, and (*S*)-ketamine on CPP scores in mice after repeated intranasal administration

In the conditioned place preference (CPP) test (Figure 4A), both (*R,S*)-ketamine and (*S*)-ketamine, but not (*R*)-ketamine, increased CPP scores, in a dose dependent manner (Figure 4). Repeated intranasal administration of (*R,S*)-ketamine (40 mg/kg), but not the low doses (10 and 20 mg/kg), significantly increased CPP scores ($F_{3,34} = 3.054$, $P = 0.042$) (Figure 4B). In contrast, repeated intranasal administration of (*R*)-ketamine (10, 20 or 40 mg/kg) did not increase CPP scores ($F_{3,36} = 0.072$, $P = 0.974$) (Figure 4C). Repeated intranasal administration of (*S*)-ketamine (20 and 40 mg/kg), but not the low dose (10 mg/kg), significantly increased CPP scores ($F_{3,36} = 14.0$, $P < 0.001$) (Figure 4D).

Collectively, the order of potencies of side effects (i.e., psychosis, abuse liability) in mice after intranasal administration was (*S*)-ketamine > (*R,S*)-ketamine > (*R*)-ketamine.




Figure 4. Effects of (*R,S*)-ketamine, (*R*)-ketamine, and (*S*)-ketamine on CPP score after repeated intranasal administration

(A): Schedule of habituation, treatment, and behavioral test. (B): Saline (0.5 ml/kg), or (*R,S*)-ketamine (10, 20, or 40 mg/kg) was administered intranasally into male mice. (C): Saline (0.5 ml/kg), or (*R*)-ketamine (10, 20, or 40 mg/kg) was administered intranasally into male mice. (D): Saline (0.5 ml/kg), or (*S*)-ketamine (10, 20, or 40 mg/kg) was administered intranasally into male mice. CPP score was measured as described in the method section. The values represent the mean \pm S.E.M. ($n = 8 - 10$). * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$ compared with saline-treated mice. N.S.: not significant. R-KT: (*R*)-ketamine. RS-KT: (*R,S*)-ketamine. S-KT: (*S*)-ketamine.

Discussion

In the present study, we compared (*R,S*)-ketamine, and its two enantiomers, in CSDS susceptible mice (for antidepressant effects) and control mice (for side effects). The order of potency of antidepressant effects after a single intranasal administration to CSDS susceptible mice is (*R*)-ketamine > (*R,S*)-ketamine > (*S*)-ketamine. Furthermore, the order of potency of side effects (i.e., psychosis and abuse liability) after intranasal administration is (*S*)-ketamine > (*R,S*)-ketamine > (*R*)-ketamine. Collectively, it is likely that (*R*)-ketamine would be a rapid-acting and sustained antidepressant without side effects compared to (*R,S*)-ketamine and (*S*)-ketamine.

In this study, we found that antidepressant effects of (*R,S*)-ketamine and its two enantiomers in CSDS susceptible mice after a single intranasal administration may be less potent than those of a single i.p. administration (Dong et al., 2017; Yang et al., 2015; 2017a; 2017b; 2018a; Zhang et al., 2015; Zhang and Hashimoto, 2019b). Lower bioavailability of intranasal administration of (*R,S*)-ketamine and its two enantiomers may contribute to lower efficacy of intranasal administration compared to i.p. administration. Interestingly, the potency of antidepressant effects of (*R,S*)-ketamine and its two enantiomers was not correlated with the potencies of these compounds at the NMDAR (Ebert et al., 1997), suggesting that NMDAR inhibition may not play a key role in the antidepressant effects of (*R,S*)-ketamine and its enantiomers. Previously, Fukumoto et al. (2017) reported that (*R,S*)-ketamine and (*R*)-ketamine, but not (*S*)-ketamine, show antidepressant effects in rats with repeated corticosterone treatments after a single i.p. administration, consistent with our current data.

Due to its serious side effects, clinical use of ketamine has remained limited (Domino, 2010; Sanacora et al., 2017; Singh et al., 2017; Zhu et al., 2016), although it has been used as an off-label antidepressant in the USA (Reardon, 2018; Wilkinson et al., 2017). In this study,

we found that locomotion after a single intranasal administration of (*R*)-ketamine is lower than those of (*R,S*)-ketamine and (*S*)-ketamine, consistent with the previous reports (Ryder et al., 1978; Yang et al., 2015) of subcutaneous or i.p. administration. Furthermore, we found that a single intranasal administration of (*R*)-ketamine did not cause PPI deficits in mice compared to (*R,S*)-ketamine and (*S*)-ketamine, consistent with the previous reports of i.p. administration (Yang et al., 2015). Interestingly, it was reported that the ED50 of (*R*)-ketamine (6.33 mg/kg) for PPI deficits in rats was higher than that of (*S*)-ketamine (2.86 mg/kg), indicating that (*S*)-ketamine disrupts PPI with 2.5-fold higher potency than (*R*)-ketamine (Halberstadt et al., 2016). Finally, we found that repeated intranasal administration of (*R*)-ketamine did not increase CPP scores in mice although (*R,S*)-ketamine and (*S*)-ketamine increased CPP scores, in a dose dependent manner, consistent with the previous reports of i.p. administration (Yang et al., 2015; 2018b). A PET study showed that a single i.v. infusion of (*S*)-ketamine (0.5 mg/kg for 40-min), but not (*R*)-ketamine (0.5 mg/kg for 40-min), produced a marked reduction of dopamine D2/3 receptor binding in conscious monkey striatum, suggesting that (*S*)-ketamine-induced dopamine release might be associated with acute psychotomimetic and dissociative side effects in humans (Hashimoto et al., 2017). Unlike (*R,S*)-ketamine and (*S*)-ketamine, it seems that intranasal infusion of (*R*)-ketamine does not appear to cause psychotomimetic effects or have abuse potential in humans, based on the lack of behavioral abnormalities (e.g., PPI deficits, CPP) observed in mice after single or repeated intranasal administration (Hashimoto, 2016a; 2016b; 2016c).

Mathisen et al. (1995) reported that the incidence of side effects (i.e., blurred vision, altered hearing, dizziness, proprioceptive disturbances, illusions) of (*S*)-ketamine (0.45 mg/kg, i.m.) treated group in patients with oral pain was higher than (*R*)-ketamine (1.8 mg/kg, i.m.) treated group, although the dose of (*S*)-ketamine (0.45 mg/kg) is lower than (*R*)-ketamine (1.8 mg/kg). Furthermore, it is also reported that experiencing illusion and alterations in

hearing, vision, and proprioception is attributable to (*S*)-ketamine's actions (Oye et al., 1992; Vollenweider et al., 1997), whereas the feelings of relaxation are associated with (*R*)-ketamine's actions (Vollenweider et al., 1997; Zanos et al., 2018). Taken all together, it seems likely that (*S*)-ketamine contributes to the acute psychotomimetic and dissociative effects of (*R,S*)-ketamine, whereas (*R*)-ketamine may not be associated with these side effects (Zanos et al., 2018).

On March 5, 2019, the US FDA approved nasal spray of (*S*)-ketamine for treatment-resistant depression (FDA 2019). Due to the risk of serious adverse outcomes from sedation and dissociation caused by administration of (*S*)-ketamine, as well as the potential for abuse and misuse of the drug, FDA said that the drug will only be available through a restricted distribution system, under a Risk Evaluation and Mitigation Strategy (REMS). Patients will self-administer (*S*)-ketamine under the supervision of a health care provider in a certified doctor's office or clinic; the nasal spray cannot be taken home (FDA 2019). Given the lack of adverse side effects of (*R*)-ketamine, it is possible that patients may take (*R*)-ketamine to their home.

In conclusion, this study suggests that the order of potency for antidepressant effects in a CSDS model after a single intranasal administration is (*R*)-ketamine > (*R,S*)-ketamine > (*S*)-ketamine. In contrast, the order of potency for side effects in mice after intranasal administration is (*S*)-ketamine > (*R,S*)-ketamine > (*R*)-ketamine. Therefore, it is likely that (*R*)-ketamine could be a safer antidepressant without side effects than (*R,S*)-ketamine and (*S*)-ketamine.

Acknowledgements

This study was supported by AMED (to K.H., JP19dm0107119). Dr. Lijia Chang was supported by the Japan China Sasakawa Medical Fellowship (Tokyo, Japan). Dr. Zhongwei Xiong (Wuhan University, China) was supported by the China Scholarship Council (China).

References

- Abdallah, C.G., Sanacora, G., Duman, R.S., Krystal, J.H., (2018). The neurobiology of depression, ketamine and rapid-acting antidepressants: Is it glutamate inhibition or activation? *Pharmacol. Ther.* 190, 148-158.
- Berman, R.M., Cappiello, A., Anand, A., Oren, D.A., Heninger, G.R., Charney, D.S., Krystal, J.H., (2000). Antidepressant effects of ketamine in depressed patients. *Biol. Psychiatry* 47, 351–354.
- Chaki, S., (2017a). Beyond ketamine: new approaches to the development of safer antidepressants. *Curr. Neuropharmacol.* 15, 963–976.
- Chaki, S., (2017b). mGlu2/3 receptor antagonists as novel antidepressants. *Trends Pharmacol. Sci.* 38, 569–580.
- Chang, L., Zhang, K., Pu, Y., Qu, Y., Wang, S.M., Xiong, Z., Shirayama, Y., Hashimoto, K., (2019). Lack of dopamine D1 receptors in the antidepressant actions of (*R*)-ketamine in a chronic social defeat stress model. *Eur. Arch. Psychiatry Clin. Neurosci.* 2019 Mar 29. doi: 10.1007/s00406-019-01012-1.
- Domino, E.F., (2010). Taming the ketamine tiger. 1965. *Anesthesiology* 113, 678–684.
- Dong, C., Zhang, J.C., Yao, W., Ren, Q., Ma, M., Yang, C., Chaki, S., Hashimoto, K., (2017). Rapid and sustained antidepressant action of the mGlu2/3 receptor antagonist MGS0039 in the social defeat stress model: Comparison with ketamine. *Int. J. Neuropsychopharmacol.* 20, 228–236.
- Duman, R.S., (2018). Ketamine and rapid-acting antidepressants: a new era in the battle against depression and suicide. *F1000Res.* 7.pii: F1000 Faculty Rev-659.
- Ebert, B., Mikkelsen, S., Thorkildsen, C., Borgbjerg, F.M., (1997). Norketamine, the main metabolite of ketamine, is a non-competitive NMDA receptor antagonist in the rat cortex and spinal cord. *Eur. J. Pharmacol.* 333, 99–104.

- FDA News Release on March 5, (2019). FDA approves new nasal spray medication for treatment-resistant depression; available only at a certified doctor's office or clinic. <https://www.fda.gov/NewsEvents/Newsroom/PressAnnouncements/ucm632761.htm>
- Fukumoto, K., Toki, H., Iijima, M., Hashihayata, T., Yamaguchi, J.I., Hashimoto, K., Chaki, S., (2017). Antidepressant Potential of (*R*)-Ketamine in Rodent Models: Comparison with (*S*)-Ketamine. *J. Pharmacol. Exp. Ther.* 361, 9–16.
- Golden, S.A., Covington III, H.E., Berton, O., Russo, S.J., (2011). A standardized protocol for repeated social defeat stress in mice. *Nat. Protoc.* 6,1183–1191.
- Gould, T.D., Zarate, C.A. Jr., Thompson, S.M., (2019). Molecular Pharmacology and Neurobiology of Rapid-Acting Antidepressants. *Annu. Rev. Pharmacol. Toxicol.* 59, 213–236.
- Grunebaum, M.F., Galfalvy, H.C., Choo, T.H., Keilp, J.G., Moitra, V.K., Parris, M.S., Marver, J.E., Burke, A.K., Milak, M.S., Sublette, M.E., Oquendo, M.A., Mann, J.J., (2018). Ketamine for rapid reduction of suicidal thoughts in major depression: a midazolam-controlled randomized clinical trial. *Am. J. Psychiatry* 175, 327–335.
- Halberstadt, A.L., Slepak, N., Hyun, J., Buell, M.R., Powell, S.B., (2016). The novel ketamine analog methoxetamine produces dissociative-like behavioral effects in rodents. *Psychopharmacology (Berl)* 233, 1215–1225.
- Hashimoto, K., (2014). The R-stereoisomer of ketamine as an alternative for ketamine for treatment-resistant major depression. *Clin. Psychopharmacol. Neurosci.* 12, 72–73.
- Hashimoto, K., (2016a). R-ketamine: a rapid-onset and sustained antidepressant without risk of brain toxicity. *Psychol. Med.* 46, 2449–2451.
- Hashimoto, K., (2016b). Ketamine's antidepressant action: beyond NMDA receptor inhibition. *Expert. Opin. Ther. Targets.* 20, 1389–1392.

- Hashimoto, K., (2016c). Detrimental side effects of repeated ketamine infusions in the brain. *Am. J. Psychiatry* 173, 1044–1045.
- Hashimoto, K., Kakiuchi, T., Ohba, H., Nishiyama, S., Tsukada, H., (2017). Reduction of dopamine D2/3 receptor binding in the striatum after a single administration of esketamine, but not R-ketamine: A PET study in conscious monkeys. *Eur. Arch. Psychiatry Clin. Neurosci.* 267, 173–176.
- Kishimoto, T., Chawla, J.M., Hagi, K., Zarate, C.A., Kane, J.M., Bauer, M., Correll, C.U., (2016). Single-dose infusion ketamine and non-ketamine N-methyl-d-aspartate receptor antagonists for unipolar and bipolar depression: a meta-analysis of efficacy, safety and time trajectories. *Psychol. Med.* 46, 1459-1472.
- Krystal, J.H., Abdallah, C.G., Sanacora, G., Charney, D.S., Duman, R.S., (2019). Ketamine: A paradigm shift for depression research and treatment. *Neuron* 101, 774-778.
- Larkin, G.L., Beautrais, A.L., (2011). A preliminary naturalistic study of low-dose ketamine for depression and suicide ideation in the emergency department. *Int. J. Neuropsychopharmacol.* 14, 1127–1131.
- Li, L., Vlisides, P.E., (2016). Ketamine: 50 years of modulating the mind. *Front. Hum. Neurosci.* 10, 612.
- Mathisen, L.C., Skjelbred, P., Skoglund, L.A., Oye, I., (1995). Effect of ketamine, an NMDA receptor inhibitor, in acute and chronic orofacial pain. *Pain* 61, 215–220.
- Matsuura, A., Fujita, Y., Iyo, M., Hashimoto, K., (2015). Effects of sodium benzoate on pre-pulse inhibition deficits and hyperlocomotion in mice after administration of phencyclidine. *Acta. Neuropsychiatr.* 27, 159–167.
- Monteggia, L.M., Zarate Jr, C., (2015). Antidepressant actions of ketamine: from molecular mechanisms to clinical practice. *Curr. Opin. Neurobiol.* 30, 139–143.

- Murrough, J.W., Abdallah, C.G., Mathew, S.J., (2017). Targeting glutamate signalling in depression: progress and prospects. *Nat. Rev. Drug Discov.* 16, 472–486.
- Murrough, J.W., Iosifescu, D.V., Chang, L.C., Al Jurdi, R.K., Green, C.E., Perez, A.M., Iqbal, S., Pillemer, S., Foulkes, A., Shah, A., Charney, D.S., Mathew, S.J., (2013). Antidepressant efficacy of ketamine in treatment-resistant major depression: a two-site randomized controlled trial. *Am. J. Psychiatry* 170, 1134–1142.
- Murrough, J.W., Soleimani, L., DeWilde, K.E., Collins, K.A., Lapidus, K.A., Iacoviello, B.M., Lener, M., Kautz, M., Kim, J., Stern, J.B., Price, R.B., Perez, A.M., Brallier, J.W., Rodriguez, G.J., Goodman, W.K., Iosifescu, D.V., Charney, D.S., (2015). Ketamine for rapid reduction of suicidal ideation: a randomized controlled trial. *Psychol. Med.* 45, 3571–3580.
- Newport, D.J., Carpenter, L.L., McDonald, W.M., Potash, J.B., Tohen, M., Nemeroff, C.B., (2015). APA Council of Research Task Force on Novel Biomarkers and Treatments. Ketamine and other NMDA antagonists: early clinical trials and possible mechanisms in depression. *Am. J. Psychiatry* 172, 950–966.
- Oye, I., Paulsen, O., Maurset, A., (1992). Effects of ketamine on sensory perception: evidence for a role of N-methyl-D-aspartate receptors. *J. Pharmacol. Exp. Ther.* 260, 1209–1213.
- Peltoniemi, M.A., Hagelberg, N.M., Olkkola, K.T., Saari, T., (2016). Ketamine: a review of clinical pharmacokinetics and pharmacodynamics in anesthesia and pain therapy. *Clin. Pharmacokinet.* 55, 1059–1077.
- Price, R.B., Nock, M.K., Charney, D.S., Mathew, S.J., (2009). Effects of intravenous ketamine on explicit and implicit measures of suicidality in treatment-resistant depression. *Biol. Psychiatry* 66, 522–526.

- Reardon, S., (2018). ‘Party drug’ turned antidepressant approaches approval. *Nat. Rev. Drug Discov.* 17, 773–775.
- Ryder, S., Way, W.L., Trevor, A.J., (1978). Comparative pharmacology of the optical isomers of ketamine in mice. *Eur. J. Pharmacol.* 49, 15–23.
- Sanacora, G., Frye, M.A., McDonald, W., Mathew, S.J., Turner, M.S., Schatzberg, A.F., Summergrad, P., Nemeroff, C.B.; American Psychiatric Association (APA) Council of Research Task Force on Novel Biomarkers and Treatments, (2017). A consensus statement on the use of ketamine in the treatment of mood disorders. *JAMA Psychiatry* 74, 399–405.
- Singh, I., Morgan, C., Curran, V., Nutt, D., Schlag, A., McShane, R., (2017). Ketamine treatment for depression: opportunities for clinical innovation and ethical foresight. *Lancet Psychiatry* 4, 419–426.
- Su, T.P., Chen, M.H., Li, C.T., Lin, W.C., Hong, C.J., Gueorguieva, R., Tu, P.C., Bai, Y.M., Cheng, C.M., Krystal, J.H., (2017). Dose-related effects of adjunctive ketamine in Taiwanese patients with treatment-resistant depression. *Neuropsychopharmacology* 42, 2482–2492.
- Tian, Z., Dong, C., Fujita, A., Fujita, Y., Hashimoto, K., (2018). Expression of heat shock protein HSP-70 in the retrosplenial cortex of rat brain after administration of (*R,S*)-ketamine and (*S*)-ketamine, but not (*R*)-ketamine. *Pharmacol. Biochem. Behav.* 172, 17–21.
- Vollenweider, F.X., Leenders, K.L., Oye, I., Hell, D., Angst, J., (1997). Differential psychopathology and patterns of cerebral glucose utilisation produced by (*S*)- and (*R*)-ketamine in healthy volunteers using positron emission tomography (PET). *Eur. Neuropsychopharmacol.* 7, 25–38.

- Wilkinson, S.T., Ballard, E.D., Bloch, M.H., Mathew, S.J., Murrrough, J.W., Feder, A., Sos, P., Wang, G., Zarate Jr, C.A., Sanacora, G., (2018). The effect of a single dose of intravenous ketamine on suicidal ideation: A systematic review and individual participant data meta-analysis. *Am. J. Psychiatry* 175, 150–158.
- Wilkinson, S.T., Toprak, M., Turner, M.S., Levine, S.P., Katz, R.B., Sanacora, G., (2017). A survey of the clinical, off-label use of ketamine as a treatment for psychiatric disorders. *Am. J. Psychiatry* 174, 695–696.
- Xiong, Z., Zhang, K., Ishima, T., Ren, Q., Chang, L., Chen, J., Hashimoto, K., (2018a). Comparison of rapid and long-lasting antidepressant effects of negative modulators of $\alpha 5$ -containing GABAA receptors and (*R*) ketamine in a chronic social defeat stress model. *Pharmacol. Biochem. Behav.* 175, 139–145.
- Xiong, Z., Zhang, K., Ishima, T., Ren, Q., Ma, M., Pu, Y., Chang, L., Chen, J., Hashimoto, K., (2018b). Lack of rapid antidepressant effects of Kir4.1 channel inhibitors in a chronic social defeat stress model: Comparison with (*R*)-ketamine. *Pharmacol. Biochem. Behav.* 176, 57–62.
- Xu, Y., Hackett, M., Carter, G., Loo, C., Gálvez, V., Glozier, N., Glue, P., Lapidus, K., McGirr, A., Somogyi, A.A., Mitchell, P.B., Rodgers, A., (2016). Effects of low-dose and very low-dose ketamine among patients with major depression: a systematic review and meta-analysis. *Int. J. Neuropsychopharmacol.* 19, pii: pyv124
- Yang, C., Han, M., Zhang, J.C., Ren, Q., Hashimoto, K., (2016). Loss of parvalbumin-immunoreactivity in mouse brain regions after repeated intermittent administration of esketamine, but not R-ketamine. *Psychiatric. Res.* 239, 281–283.
- Yang, C., Kobayashi, S., Nakao, K., Dong, C., Han, M., Qu, Y., Ren, Q., Zhang, J.C., Ma, M., Toki, H., Yamaguchi, J.I., Chaki, S., Shirayama, Y., Nakazawa, K., Manabe, T.,

- Hashimoto, K., (2018b). AMPA receptor activation-independent antidepressant actions of ketamine metabolite (S)-norketamine. *Biol. Psychiatry* 84, 591–600.
- Yang, C., Qu, Y., Abe, M., Nozawa, D., Chaki, S., Hashimoto, K., (2017a). (*R*)-ketamine shows greater potency and longer lasting antidepressant effects than its metabolite (2R,6R)-hydroxynorketamine. *Biol. Psychiatry* 82, e43–e44.
- Yang, C., Qu, Y., Fujita, Y., Ren, Q., Ma, M., Dong, C., Hashimoto, K., (2017b). Possible role of the gut microbiota-brain axis in the antidepressant effects of (*R*)-ketamine in a social defeat stress model. *Transl. Psychiatry* 7, 1294.
- Yang, C., Ren, Q., Qu, Y., Zhang, J.C., Ma, M., Dong, C., Hashimoto, K., (2018a). Mechanistic target of rapamycin-independent antidepressant effects of (*R*)-ketamine in a social defeat stress model. *Biol. Psychiatry* 83, 18–28.
- Yang, C., Shirayama, Y., Zhang, J.C., Ren, Q., Yao, W., Ma, M., Dong, C., Hashimoto, K., (2015). *R*-ketamine: a rapid-onset and sustained antidepressant without psychotomimetic side effects. *Transl. Psychiatry* 5, e632.
- Zanos, P., Moaddel, R., Morris, P.J., Georgiou, P., Fischell, J., Elmer, G.I., Alkondon, M., Yuan, P., Pribut, H.J., Singh, N.S., Dossou, K.S., Fang, Y., Huang, X.P., Mayo, C.L., Wainer, I.W., Albuquerque, E.X., Thompson, S.M., Thomas, C.J., Zarate Jr, C.A., Gould, T.D., (2016). NMDAR inhibition-independent antidepressant actions of ketamine metabolites. *Nature* 533, 481–486.
- Zanos, P., Moaddel, R., Morris, P.J., Riggs, L.M., Highland, J.N., Georgiou, P., Pereira, E.F.R., Albuquerque, E.X., Thomas, C.J., Zarate Jr, C.A., Gould, T.D., (2018). Ketamine and ketamine metabolite pharmacology: insights into therapeutic mechanisms. *Pharmacol. Rev.* 70, 621–660.

- Zarate, C.A., Singh, J.B., Carlson, P.J., Brutsche, N.E., Ameli, R., Luckenbaugh, D.A., Charney, D.S., Manji, H.K., (2006). A randomized trial of an N-methyl-D-aspartate antagonist in treatment-resistant major depression. *Arch. Gen. Psychiatry* 63, 856–864.
- Zhang, K., Hashimoto, K., (2019a). An update on ketamine and its two enantiomers as rapid-acting antidepressants. *Expert Rev. Neurother.* 19, 83–92.
- Zhang, K., Hashimoto, K., (2019b). Lack of opioid system in the antidepressant actions of ketamine. *Biol. Psychiatry* 85, e25–e27.
- Zhang, J.C., Li, S.X., Hashimoto, K., (2014). *R*(-)-ketamine shows greater potency and longer lasting antidepressant effects than *S*(+)-ketamine. *Pharmacol. Biochem. Behav.* 116, 137–141.
- Zhang, K., Toki, H., Fujita, Y., Ma, M., Chang, L., Qu, Y., Harada, S., Nemoto, T., Mizuno-Yasuhira, A., Yamaguchi, J.I., Chaki, S., Hashimoto, K., (2018). Lack of deuterium isotope effects in the antidepressant effects of (*R*)-ketamine in a chronic social defeat stress model. *Psychopharmacology (Berl)* 235, 3177–3185.
- Zhang, J.C., Yao, W., Dong, C., Yang, C., Ren, Q., Ma, M., Han, M., Hashimoto, K., (2015). Comparison of ketamine, 7,8-dihydroxyflavone, and ANA-12 antidepressant effects in the social defeat stress model of depression. *Psychopharmacology (Berl)* 232, 4325–4335.
- Zhu, W., Ding, Z., Zhang, Y., Shi, J., Hashimoto, K., Lu, L., (2016). Risks associated with misuse of ketamine as a rapid-acting antidepressant. *Neurosci. Bull.* 32, 557–564.

Pharmacology Biochemistry and Behavior 181: 53-59

2019年4月26日公表済

DOI: 10.1016/j.pbb.2019.04.008